arXiv:1209.3571v2 [math.AG] 10 Oct 2012

ON THE SLOPE CONJECTURE FOR THE FOURGONAL LOCUS IN $\overline{\mathcal{M}}_g$

BEORCHIA VALENTINA AND ZUCCONI FRANCESCO

ABSTRACT. We prove the Harris - Stankova slope conjecture for some families $\mathcal{B}$ of curves contained in $\overline{\mathcal{M}}^1_{g,4}$ such that the closure $\overline{\cup\mathcal{B}}$ is $\overline{\mathcal{M}}^1_{g,4}$. We also refine Harris - Stankova slope conjecture for trigonal curves of odd genus.

CONTENTS

1. Introduction 1
2. Fibrations factorizing through finite covers of a ruled surface 5
2.1. The trigonal case 7
2.2. The fourgonal case 9
3. Slope of general semistable fibrations in trigonal and fourgonal curves. 12
3.1. The trigonal case 14
3.2. The fourgonal case 16
References 19

1. INTRODUCTION

The geometry of the moduli space $\overline{\mathcal{M}}_g$ is strictly related to the properties of the effective cone $\mathbb{E} \subset Pic(\overline{\mathcal{M}}_g) \otimes \mathbb{R}$ and of the slope function $s\colon \mathbb{E} \to \mathbb{R} \cup \{\infty\}$ of effective divisors on $\overline{\mathcal{M}}_g$, as defined by Harris and Morrison in [HMo]

$$s(D) := \inf\{\frac{a}{b} : a, b > 0 \text{ such that } a\lambda - b\delta - D \equiv \sum_{i=0}^{[g/2]} c_i\delta_i, \text{ where } c_i \geq 0\},$$

where λ denotes the class of the Hodge line bundle, $\delta_0, \dots, \delta_{[g/2]}$ the boundary divisor classes corresponding to singular stable curves and $\delta := \delta_0 + \cdots + \delta_{[g/2]}$ the total boundary.

Then one can define the slope of the moduli space as

$$s(\overline{\mathcal{M}}_g) := \inf\{s(D) : D \in \mathbb{E}\}.$$

The number $s(\overline{\mathcal{M}}_g)$ is a meaningful invariant of $\overline{\mathcal{M}}_g$ and it encodes several information on the geometry of $\overline{\mathcal{M}}_g$, see: [EH], [Fa], [FP], [HM], [HMo]. For instance, the Harris-Mumford theorem [HM] on the Kodaira dimension of $\overline{\mathcal{M}}_g$, is equivalent to the inequality $s(\overline{\mathcal{M}}_g) < s(K_{\overline{\mathcal{M}}_g}) = \frac{13}{2}$, for $g \geq 24$. Harris and Morrison conjectured [HMo] that the inequality $s(\overline{\mathcal{M}}_g) \geq 6 + \frac{12}{g+1}$ holds, equality being attained only for the classical Brill-Noether divisors. Counterexamples were provided in [FP] for infinitely many g though all of these have slope greater than 6. On the other hand, all the methods (cf. [HMo, P]) for bounding $s(\overline{\mathcal{M}}_g)$ from below for large g, yield only bounds that tend to zero with g.

1991 *Mathematics Subject Classification.* Primary 14D22, 14H51; Secondary 14H10.

Key words and phrases. chern classes, moduli of curves, fibration, gonality.

Recently Chen, Farkas and Morrison in [CFM] posed the following natural question:

Problem 1.1. Does the limit $s_\infty := \lim_{g\to\infty} s(\overline{\mathcal{M}}_g)$ exist, and, if so, what is its value?

A possible approach to such a problem is the following: find a Zariski dense collection $\mathcal{B}$ of 1-cycles B_μ, so that any effective divisor must intersect one of these curves non-negatively, obtaining the bound $s(\overline{\mathcal{M}}_g) \geq \inf_\mu s(B_\mu)$, where $s(B_\mu) = \left(\frac{B_\mu \cdot \delta}{B_\mu \cdot \lambda}\right)$.

To such a circle of ideas belongs the study of the bounds on the slope on the moduli spaces of n-gonal curves $\overline{\mathcal{M}}^1_{g,n}$. A conjecture of Harris and Stankova, see [S] Conjecture 13.3, asserts that if $B \subset \overline{\mathcal{M}}^1_{g,n}$ is any 1-dimensional family with smooth generic member and not lying in a codimension one subvariety of $\overline{\mathcal{M}}^1_{g,n}$, then:

$$s(B) \leq \left(6 + \frac{2}{n-1}\right) + \frac{2n}{g}.$$

The aim of this paper is to construct families of n-gonal curves containing the general element of the families of curves constructed in [HMo] and such that the Harris-Stankova conjecture holds for them.

To explain our work, let us consider a curve $B \subset \overline{\mathcal{M}}^1_{g,n}$ and let us perform a semistable reduction $f\colon S \to \overline{B}$, see [AW]. Recall that the notion of slope $s(B)$ for a curve $B \subset \overline{\mathcal{M}}_g$ is related to the slope $s(f)$ of $f\colon S \to \overline{B}$. If $\omega_f := \omega_S \otimes f^\star \omega^{-1}_{\overline{B}}$ denotes the relative canonical bundle, $K_f = K_S - f^\star K_{\overline{B}}$ is the associated divisor and $\chi_f = \deg f_\star \omega_f$ denotes the relative Euler characteristic, Xiao (cf. [X1]) defines the slope of f as

$$s(f) = \frac{K_f^2}{\chi_f}.$$

The slope $s(f)$ of semistable fibrations is invariant under base change (cf. [Sz]).

We have the equality $\lambda \cdot B = \chi_f$, and Grothendieck-Riemann-Roch Theorem applied to the projection from the universal curve to the subscheme of canonical stable curves of genus g gives the relation (see [M]) $12(\lambda \cdot B) = K_f^2 + \delta \cdot B$. Then

$$s(f) = 12 - s(B) \in (0, 12].$$

Hence the slope conjecture for a general $B \subset \overline{\mathcal{M}}^1_{g,n}$ can be rewritten as $s(f) \geq 6 - \frac{2}{n-1} - \frac{2n}{g} =: \mathcal{F}_n(g)$.

The first result relating the gonality of the general fiber and the slope was given by Cornalba-Harris in [CH] and Xiao in [X2]:

Theorem 1.2. *(Cornalba-Harris, Xiao). If $g \geq 2$ and f is not locally trivial, then $s(f) \geq 4 - \frac{4}{g}$. Moreover, $s(f) = 4 - \frac{4}{g}$ if and only if the general fiber is a hyperelliptic curve.*

On the other hand, if B passes through a general point of $\overline{\mathcal{M}}_g$, Eisenbud - Harris [EH] and Konno [K2] gave the following better bound:

Theorem 1.3. *(Eisenbud-Harris, Konno). Assume that the general fibre of a fibration in semistable curves is general in its moduli space. Then $s(f) \geq 6\,\frac{(g-1)}{(g+1)}$.*

The families not satisfying the last inequality are contained in the closure of the locus of n-gonal curves $\overline{\mathcal{M}}^1_{g,n} \subset \overline{\mathcal{M}}_g$ for $n = \lfloor \frac{g+3}{2} \rfloor - 1$.

A first result toward the slope conjecture has been given by [S]:

Theorem 1.4. *(Stankova) The slope of any fibration in Deligne-Mumford semistable trigonal curves satisfies*

$$s(f) \geq \frac{24(g-1)}{5g+1},$$

with equality if and only if all fibers are irreducible, S is a triple cover of a ruled surface Y over B, the ramification R divisor of the covering $\rho\colon S \to Y$ satisfies $R \equiv \frac{1}{3}\rho^\star\rho_\star R$, and the general fiber of $f\colon S \to B$ is a trigonal curve of Maroni invariant $m \geq 2$.

Furthermore, if g is even, if the general fiber has Maroni invariant 0 and a condition on the singularities of the singular fibers is satisfied, then the slope is bounded by:

$$s(f) \geq 5 - \frac{6}{g}.$$

Barja and Stoppino [BS] proved that the last bound for g even holds without the hypotheses of semistability and without the assumptions on the singularities by using the Cornalba-Harris inequality. Another proof of such a result in even genus case has been given recently by Fedorchuk and Jensen in [FJ].

Some particular examples of fourgonal fibrations are given by double covers of hyperelliptic fibrations. Concerning these cases we have the following results [BZ, B]:

Theorem 1.5. *Barja-Zucconi: Let $f\colon S \to B$ be a genus g, relatively minimal, non isotrivial fibration. Assume f is a double cover fibration of a fibration of genus γ. Then, if $g \geq 4\gamma + 1$ we have*

$$s(f) \geq 4 + 4\frac{(\gamma-1)(g-4\gamma-1)}{(g-4\gamma-1)(g-\gamma)+2(g-1)\gamma^2}.$$

Recently, such a result has been sharpened by Cornalba and Stoppino in [CS]:

Theorem 1.6. *Cornalba - Stoppino: Let $f : X \to B$ be a double fibration of type (g, γ), that is a relatively minimal genus g fibered surface such that there is a degree 2 morphism from the general fiber of f to a smooth curve of genus γ.*

Then

$$s(f) \geq 4 + 4\frac{(\gamma-1)}{(g-\gamma)}.$$

In the present paper we investigate the cases $n = 3$ and $g \geq 5$ and $n = 4$ and $g \geq 10$.

Let $B \subset \overline{\mathcal{M}}^1_{g,3}, \overline{\mathcal{M}}^1_{g,4}$ be a curve not contained in the boundary. By the semistable reduction Theorem, up to base change over the normalization of B, we can construct a semistable fibration $f\colon S \to \overline{B}$ such that the closure of the image of the induced moduli map $\overline{B} \dashrightarrow \overline{\mathcal{M}}_g$ is B, and $f : S \to \overline{B}$ factorizes through a rational map $\rho\colon S \dashrightarrow Y$ to a ruled surface $\pi_B\colon Y \to \overline{B}$ (see, for instance, [HMo]). We next resolve the map ρ, and we get a *generically finite* morphism $\pi\colon X \to Y$. The case where $\pi\colon X \to Y$ is finite is treated in Section 2. To deal with the difficult case where $\pi\colon X \to Y$ is not finite we say that a curve $B \subset \overline{\mathcal{M}}^1_{g,n}$ is an HM-curve if $\pi\colon X \to Y$ is a finite morphism over a neighbourhood of any point of the branch locus $B(\pi)$, which is not a simple node; see: Definition 3.2. If $B \subset \overline{\mathcal{M}}^1_{g,3}, \overline{\mathcal{M}}^1_{g,4}$ is an HM-curve the analysis of the locus where the morphism $\pi\colon X \to Y$ is not finite is essentially given by Harris and Morrison in [HMo, pages 343-347] and by their study we see that the family $\mathcal{B}$ of HM-curves is such that $\overline{\cup\mathcal{B}}$ is $\overline{\mathcal{M}}^1_{g,n}$ for every n.

Next, instead of pursuing the approach explained in [HMo], we consider the Stein factorization of $\pi\colon X \to Y$ to get a finite flat morphism $h\colon \widehat{X} \to Y$ between a normal surface $\widehat{X}$ and the ruled surface Y. Then we perform a partial resolution of $h\colon \widehat{X} \to Y$ to obtain a Gorenstein cover $\widetilde{\rho}\colon \widetilde{X} \to \widetilde{Y}$ where $\widetilde{Y}$ is a suitable blow-up of the ruled surface Y. Since we consider only HM-curves and the cover has degree at most four, we can control how the invariants change along the process. In particular, by the theory of Gorenstein covers, we are able to express the invariants of the induced fibration $\widetilde{X} \to \overline{B}$ in terms of the Chern classes of the push-forward $\widetilde{\rho}_\star\omega_{\widetilde{X}/\widetilde{Y}}$ of the relative canonical bundle.

In the trigonal case, we recover Theorem 1.4 and the missing case of a fibration in general trigonal curves of odd genus, with the assumptions that B does not intersect the codimension two locus of curves with Maroni invariant $m \geq 3$ in $\overline{\mathcal{M}}^1_{g,3}$. The crucial result we use is Viehweg Weak Positivity Theorem [V, 3.4] for $\widetilde{\rho}_\star\omega_{\widetilde{X}/\widetilde{Y}}$. We prove:

Theorem 1.7. *If $B \subset \overline{\mathcal{M}}^1_{g,3}$ with $g \geq 5$ is an HM-curve, with general point corresponding to a general trigonal curve of odd genus $g \geq 11$, and assume that B does not intersect the codimension two locus of curves with Maroni invariant $m \geq 3$. Then*

$$s(f) \geq 5 - \frac{8}{g+1}.$$

Remark. We point out that we received a private communication by A. Deopurkar and A. Patel, who constructed a family of rational curves in $\overline{\mathcal{M}}^1_{g,3}$ with g odd and dense union, whose slope is slightly higher than our bound. Their fibrations factorize through a finite Gorenstein cover $\rho : S \to \mathbb{P}^1 \times \mathbb{P}^1$ with quotient vector bundle $(\rho_\star\omega_{S/\mathbb{P}^1\times\mathbb{P}^1})/\mathcal{O}_{\mathbb{P}^1\times\mathbb{P}^1})$ very ample and uniform.

Our technique allows to treat also the fourgonal case for HM-curves. We obtain a similar description to the trigonal case.

Theorem 1.8. *Let $B \subset \overline{\mathcal{M}}^1_{g,4}$ with $g \geq 10$ be a HM-curve, and let $f : S \to \overline{B}$ be a semistable reduction. Then:*

(1) if the general fiber of f is a general fourgonal curve of odd genus $g \geq 10$, then

$$s(f) \geq \frac{16}{3} - \frac{16}{3(3g+1)};$$

in particular, the slope conjecture is true for an HM-curve in $\overline{\mathcal{M}}^1_{g,4}$ for g odd.

If moreover the vector bundle $\widetilde{\rho}_\star\mathcal{I}_{\widetilde{X},\mathbb{P}(\widetilde{\mathcal{E}})}(2)$ arising from the associated Gorenstein cover $\widetilde{\rho} : \widetilde{X} \to \widetilde{Y}$ is weakly positive and uniform, it holds:

(2) if the general fiber of f is a general fourgonal curve of even genus, then

$$s(f) \geq \frac{16}{3} - \frac{8}{g} = \mathcal{F}_4(g);$$

(3) if for the general fiber of f the gonal degree four cover does not factorize, then

$$s(f) \geq \frac{24(g-1)}{(5g+3)},$$

with equality if and only if f factorizes through a finite degree four cover ρ of a ruled surface whose ramification divisor R satisfies $R \equiv \frac{1}{4}\rho^\star\rho_\star R$;

(4) if for the general fiber of f the gonal degree four covering factorizes through a double cover of a hyperelliptic curve of genus $\gamma < \frac{(g-3)}{6}$, then

$$s(f) \geq 4 + 4\frac{(\gamma-1)}{(g-\gamma)}.$$

In particular, the slope conjecture is true for an HM-curve under our assumptions.

Remark. The uniformity condition on $\widetilde{\rho}_\star \mathcal{I}_{\widetilde{X},\mathbb{P}(\widetilde{\mathcal{E}})}(2)$ is open, as Brundu and Sacchiero have shown in [BSa] that if C is a fourgonal curve and (α, β) is the splitting type of the analogous rank two vector bundle, the dimension of the varieties parametrizing fourgonal curves with fixed splitting type drop by two if β increases by one.

The weak positivity condition is satisfied, for instance, by a large class of complete intersections of two very ample divisors in a projective bundle over a ruled surface.

Remark. Our results confirm the general belief, that fibrations with low slope $s(f)$ should occur when the fibers are special in their moduli space.

Acknowledgment. This research has been supported by MIUR funds, PRIN project *Geometria delle varietà algebriche e dei loro spazi di moduli* (2008).

The authors are grateful to M. Barja, L. Stoppino and E. Tenni for fruitful discussion had in September 2010, when they visited the Departement of Mathematics and Informatics in Trieste. The authors thank also P. Hawking, M. Manetti, A. Verra and A. Alzati.

Finally, the authors are grateful to A. Deopurkar and A. Patel for their remarks on the earlier version of the paper, and for the explanation of their interesting examples.

2. FIBRATIONS FACTORIZING THROUGH FINITE COVERS OF A RULED SURFACE

In this section we assume that $f : S \to B$ is a fibration, which factorizes through a degree n cover $\rho : S \to Y$, where $\pi_B : Y \to B$ is a ruled surface over the smooth genus b curve B. We shall denote by

$$\mathcal{E}^\vee \cong coker\ \rho^\#,$$

where $\rho^\# : \mathcal{O}_Y \to \rho_\star \mathcal{O}_S$ is the trace map, and by R the ramification divisor. The following result is well-known, see for instance [CE].

Theorem 2.1. *Let Y be an integral surface and let $\rho : S \to Y$ be a Gorenstein cover of degree $n \geq 3$. There exists a unique $\mathbb{P}^{n-2}$-bundle $\pi_Y : \mathbb{P} \to Y$ and an embedding $i : S \to \mathbb{P}$ such that $\rho = \pi_Y \circ i$. Moreover $\mathbb{P} \cong \mathbb{P}(\mathcal{E})$ and the ramification divisor R satisfies:*

$$\mathcal{O}_S(R) \cong i^\star \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1).$$

The Grothendieck-Riemann-Roch Theorem applied to the morphism ρ gives some first relations between the invariants of S, Y and $\mathcal{E}$. We denote by $A(Y)$ the Chow ring of Y and by $\equiv$ the numerical equivalence.

Lemma 2.2. *Let Y be an integral surface. Let $\rho : S \to Y$ be a flat cover of degree n, and let $\rho_\star \mathcal{O}_S \cong \mathcal{O}_Y \oplus \mathcal{E}^\vee$. Then in $A(Y) \otimes \mathbb{Q}$ we have*

1. $\rho_\star R \equiv 2c_1(\mathcal{E})$,
2. $\chi(\mathcal{O}_S) = n\chi(\mathcal{O}_Y) + \frac{1}{2}c_1(\mathcal{E}) \cdot K_Y + \frac{1}{2}c_1(\mathcal{E})^2 - c_2(\mathcal{E})$;
3. $c_1(\rho_\star \mathcal{O}_S(2R)) \equiv 3c_1(\mathcal{E})$,
4. $c_2(\rho_\star \mathcal{O}_S(2R)) = 4c_1(\mathcal{E})^2 + c_2(\mathcal{E}) - R^2$.

Proof. The result follows from Grothendieck-Riemann-Roch Theorem applied to the morphism $\rho : S \to Y$ and the sheaves $\mathcal{O}_S$ and $\mathcal{O}_S(2R)$, respectively. The first relation is well known. We have

$$\mathrm{ch}(\rho_! \mathcal{O}_S) \cdot \mathrm{td}\mathcal{T}_Y = \rho_\star(\mathrm{ch}\mathcal{O}_S \cdot \mathrm{td}\mathcal{T}_S),$$

which gives

$$\mathrm{ch}(\rho_\star \mathcal{O}_S) \cdot (1 - \frac{1}{2}K_Y + \chi(\mathcal{O}_Y)) = \rho_\star(1 - \frac{1}{2}K_S + \chi(\mathcal{O}_S)),$$

that is

$$(n - c_1(\mathcal{E}) + \frac{1}{2}(c_1^2(\mathcal{E}) - 2c_2(\mathcal{E})) \cdot (1 - \frac{1}{2}K_Y + \chi(\mathcal{O}_Y)) = \rho_\star(1 - \frac{1}{2}K_S + \chi(\mathcal{O}_S)).$$

The divisorial part satisfies

$$-c_1(\mathcal{E}) = -\frac{1}{2}(\rho_\star K_S - 4K_Y).$$

As $K_S \sim \rho^\star K_Y + R$, we have $\rho_\star K_S \equiv 4K_Y + \rho_\star R$ and the first claim follows.

The equality between the codimension two cycles gives formula (2).

Formulae (3), (4) follow from the relation

$$\mathrm{ch}(\rho_\star \mathcal{O}_S(2R)) \cdot \mathrm{td}\mathcal{T}_Y = \rho_\star(\mathrm{ch}\mathcal{O}_S(2R) \cdot \mathrm{td}\mathcal{T}_S),$$

which implies

$$(n + c_1(\rho_\star \mathcal{O}_S(2R)) + \frac{1}{2}(c_1^2(\rho_\star \mathcal{O}_S(2R)) - 2c_2(\rho_\star \mathcal{O}_S(2R))) \cdot (1 - \frac{1}{2}K_Y + \chi(\mathcal{O}_Y)) =$$

$$= \rho_\star(1 + 2R + 2R^2) \cdot (1 - \frac{1}{2}K_S + \chi(\mathcal{O}_S))).$$

Finally we apply (1) and (2). ☐

Lemma 2.3. *Let $\rho\colon S \to Y$ be a Gorenstein degree n cover of a ruled surface $\pi_B\colon Y \to B$ over a smooth curve B. Let $T_Y \subset Y$ be any section, let F_y be the fiber over a point $y \in B$ and let T_0 be the $\mathbb{Q}$-divisor on Y given by $T_0 = T_Y - \frac{1}{2}T_Y^2 F_y$. Then*

$$c_1(\mathcal{E}) \equiv (g + n - 1)T_0 + \left(\frac{c_1(\mathcal{E})^2}{2(g + n - 1)}\right) F_y.$$

Proof. Set $\mathcal{E}_y = \mathcal{E} \otimes \mathcal{O}_{F_y}$. As T_0 and F_y form a $\mathbb{Q}$-basis of $Pic(Y)$, we can express $c_1(\mathcal{E})$ as:

$$c_1(\mathcal{E}) \equiv \deg c_1(\mathcal{E}_y) T_0 + \delta F_y$$

for some $\delta \in \mathbb{Q}$.

We claim that $\deg c_1(\mathcal{E}_y) = g + n - 1$.

Indeed, by Lemma 2.2 $\rho_\star R = 2c_1(\mathcal{E})$, and by costruction $\rho^* F_y = C_y$. Hence, by projection formula, we have $\deg c_1(\mathcal{E}_y) = \frac{1}{2} R \cdot C_y$. Since the general fiber C_y is transversal to the ramification divisor R, then the restriction $R_{|C_y}$ gives the full ramification locus of the degree n covering $\rho_y\colon C_y \to \mathbb{P}^1$. In particular the claim follows by Riemann-Hurwitz formula applied to $\rho_y : C_y \to \mathbb{P}^1$.

Finally, by taking into account that $T_0^2 = 0,\ F_y^2 = 0,\ T_0 \cdot F_y = 1$, we get $c_1(\mathcal{E})^2 = 2(g+n-1)\delta$. ☐

Proposition 2.4. *Let $f : S \to B$ be a fibration which factorizes through a Gorenstein degree n cover of a ruled surface. Then*

$$s(f) = \frac{R^2 - \frac{4}{(g+n-1)} c_1(\mathcal{E})^2}{\frac{(g+n-2)}{2(g+n-1)} c_1(\mathcal{E})^2 - c_2(\mathcal{E})}.$$

Proof. Recall that Grothendieck-Riemann-Roch Theorem applied to the fibration $f : S \to B$ yields $K_f^2 = K_S^2 - 8(g-1)(b-1)$, $\chi_f = \chi(\mathcal{O}_S) - (g-1)(b-1)$.

By Lemma 2.2 we can write:

$$K_f^2 = (R + \rho^\star K_Y)^2 - 8(g-1)(b-1) = R^2 + 4c_1(\mathcal{E}) \cdot K_Y + nK_Y^2 - 8(g-1)(b-1).$$

In the $\mathbb{Q}$-basis T_0, F_y we have $K_Y \equiv -2T_0 + \pi_B^\star K_B$, so $K_Y^2 = -8(b-1)$ and $c_1(\mathcal{E}) \cdot K_Y = -\frac{c_1(\mathcal{E})^2}{(g+n-1)} + 2(g+n-1)(b-1)$ since Lemma 2.3. Hence the computation of K_f^2 easily follows.

To determine χ_f we apply Lemma 2.2 (2) and the relations above taking into account that $\chi(\mathcal{O}_Y) = 1 - b$ since Y is ruled over B. ☐

We conclude with an upper bound on R^2, which follows from the Algebraic Index Theorem.

Lemma 2.5. *Let $\rho : S \to Y$ be a degree n Gorenstein cover, with S a normal surface, and let $\rho_\star \mathcal{O}_S \cong \mathcal{O}_Y \oplus \mathcal{E}^\vee$. Then*

$$R^2 \leq \frac{4}{n} c_1(\mathcal{E})^2.$$

Proof. By the Algebraic Index Theorem on a normal projective surface the signature of the intersection pairing on $\mathrm{Num}(S) \otimes \mathbb{R}$ is $(1, m-1)$, where m is the rank of $\mathrm{Num}(S)$.

Consider the subvector space V of $\mathrm{Num}(S) \otimes \mathbb{R}$ generated by R and $\rho^\star c_1(\mathcal{E})$. If it is 1-dimensional, then $R \equiv q \rho^\star c_1(\mathcal{E})$ for some $q \in \mathbb{R}$; as $\rho_\star R \equiv 2c_1(\mathcal{E})$, we have $q = 2/n$, so $R^2 = \frac{4}{n} c_1(\mathcal{E})^2$.

Now assume that $V := \mathbb{R}\langle R \rangle \oplus \mathbb{R}\langle \rho^\star c_1(\mathcal{E}) \rangle$ is 2-dimensional. Since R gives an embedding then intersection pairing on such a subgroup has signature $(1, -1)$.

So the determinant of the matrix

$$\begin{pmatrix} R^2 & R \cdot \rho^\star c_1(\mathcal{E}) \\ R \cdot \rho^\star c_1(\mathcal{E}) & (\rho^\star c_1(\mathcal{E}))^2 \end{pmatrix}$$

is negative, that is $R^2(nc_1(\mathcal{E})^2) \leq (R \cdot \rho^\star c_1(\mathcal{E}))^2$. As $\rho_\star R \equiv 2c_1(\mathcal{E})$, the claim follows. □

The idea now is to determine some lower bounds on $c_2(\mathcal{E})$ and on R^2 in the trigonal and fourgonal case.

2.1. **The trigonal case.**

Now we treat the case $n = 3$. We have:

Proposition 2.6. *Let $f : S \to B$ be a fibration with general fiber a trigonal curve, and assume that it factorizes through a Gorenstien degree three cover $\rho : S \to Y$ of a ruled surface $\pi_B \colon Y \to B$. Then*

$$R^2 = 2c_1(\mathcal{E})^2 - 3c_2(\mathcal{E}). \tag{2.1}$$

In particular

$$s(f) = \frac{\frac{2g}{(g+2)} c_1(\mathcal{E})^2 - 3c_2(\mathcal{E})}{\frac{(g+1)}{2(g+2)} c_1(\mathcal{E})^2 - c_2(\mathcal{E})}. \tag{2.2}$$

Proof. As the fibers of ρ are all Gorenstein zero-dimensional schemes of degree three, the sheaves $\rho_\star \mathcal{I}_{S,\mathbb{P}(\mathcal{E})}(2)$ and $R^1\rho_\star \mathcal{I}_{S,\mathbb{P}(\mathcal{E})}(2)$ are locally free, and since they are generically zero, they are zero. So the push forward of the exact sequence

$$0 \to \mathcal{I}_{S,\mathbb{P}(\mathcal{H})}(2) \to \mathcal{O}_{\mathbb{P}(\mathcal{E})}(2) \to \mathcal{O}_S(2R) \to 0$$

yields an isomorphism

$$Sym^2(\mathcal{E}) \cong \rho_\star \mathcal{O}_S(2R).$$

By taking into account that $c_1(Sym^2\mathcal{E}) \equiv 3c_1(\mathcal{E})$, $c_2(Sym^2(\mathcal{E})) \equiv 4c_2(\mathcal{E}) + 2c_1(\mathcal{E})^2$, the equation $c_2(Sym^2(\mathcal{E})) = c_2(\rho_\star \mathcal{O}_S(2R))$ gives the statement since Lemma 2.2.

□

We finally bound $c_2(\mathcal{E})$. When the general fiber is any trigonal curve, we apply Lemma 2.5.

When the general fiber is a general trigonal curve, the gonal bundle is balanced and we are able to bound $c_2(\mathcal{E})$ using Theorem 2.8.

Proposition 2.7. *Let $f : S \to B$ be a fibration with general fiber a trigonal curve, and assume that f factorizes through a Gorenstien degree three cover $\rho : S \to Y$. Then*

(1) $c_2(\mathcal{E}) \geq \frac{2}{9} c_1(\mathcal{E})^2$;

(2) $c_2(\mathcal{E}) \geq \frac{1}{4} c_1(\mathcal{E})^2$ *if the general fiber of f is a general trigonal curve of even genus g;*

(3) *$c_2(\mathcal{E}) > \frac{(g+1)}{4(g+2)} c_1(\mathcal{E})^2$ if the general fiber of f is a general trigonal curve of odd genus g, and $\mathcal{E}$ is uniform on Y.*

Proof. The first two inequalities can be found also in [S].

The first statement follows from Lemma 2.5 and the equality $R^2 = 2c_1(\mathcal{E})^2 - 3c_2(\mathcal{E})$ of Proposition 2.6. The other two statements follow from Theorem 2.8 below, taking into account that for a general trigonal curve the Maroni invariant is zero in even genus case, so the general splitting type of the restriction of $\mathcal{E}$ to a fiber of Y is (α, β) with $\alpha = \beta = \frac{(g+2)}{2}$ and $\mathcal{E}$ turns out to be Bogomolov semistable.

In the odd genus case, the general Maroni invariant is 1, so $\alpha = \beta - 1 = \frac{(g+1)}{2}$. Moreover, as $\rho_\star \omega S/Y$ is weakly positive by Viehweg's Theorem [V, 3.4], as by relative duality we have $\rho_\star \omega S/Y \cong (\rho_\star \mathcal{O}_S)^\vee$ and as the trace sequence splits, the rank two vector bundle is a direct summand of $\rho_\star \omega S/Y$, so it is weakly positive. ☐

Theorem 2.8. *Let $\mathcal{G}$ be a rank two vector bundle on a ruled surface $\pi_B : Y \to B$ over a smooth curve B, and let (α, β), $0 < \alpha \le \beta$ be the general splitting type of the restriction to a fiber of Y. Then*

(1) *$c_2(\mathcal{G}) \ge \frac{1}{4} c_1(\mathcal{G})^2$ if $\alpha = \beta$;*

(2) *$c_2(\mathcal{G}) > \frac{\alpha}{2(\alpha+\beta)} c_1(\mathcal{G})^2$ if $\alpha < \beta$ and $\mathcal{G}$ is weakly positive and uniform on Y.*

Proof. We shall follow [B].

Assume first $\alpha < \beta$ and G weakly positive and uniform. Let $T_Y \subset Y$ be a section of $\pi_B \colon Y \to B$ with $T_Y^2 \le 0$. The locally free sheaf $\pi_{B\star}\mathcal{G}(-\beta T_Y)$ on B has rank one and we set

$$\pi_{B\star}\mathcal{G}(-\beta T_Y) = \mathcal{O}_B(L)$$

for some divisor L on B. Moreover, the natural map $\pi_B^\star \mathcal{O}_B(L) \to \mathcal{G}(-\beta T_Y)$ is generically injective, hence it is an injective map of sheaves. Then we can consider the following exact sequence:

$$0 \to \mathcal{O}_Y(\pi_B^\star L) \to \mathcal{G}(-\beta T_Y) \to \mathcal{O}_Y(D) \otimes \mathcal{I}_Z \to 0,$$

where $Z \subseteq Y$ is an effective zero-dimensional cycle such that $\pi_B^\star \pi_{B\star} Z$ is the jumping lines divisor.

As $\mathcal{G}$ is uniform, we have $Z = \emptyset$. Then we have the sequence

$$0 \to \mathcal{O}_Y(\beta T_Y + \pi_B^\star L) \to \mathcal{G} \to \mathcal{O}_Y(\alpha T_Y + \pi_B^\star M) \to 0,$$

and the Chern classes satisfy

$$c_2(\mathcal{G}) = (\beta T_Y + \pi_B^\star L) \cdot (\alpha T_Y + \pi_B^\star M) = \alpha\beta T_Y^2 + \alpha \deg(\pi_B^\star L) + \beta \deg(\pi_B^\star M).$$

Next we express $c_1(\mathcal{G})$ in the following way:

$$c_1(\mathcal{G}) \equiv (\alpha+\beta) T_Y + \frac{(c_1(\mathcal{G})^2 - (\alpha+\beta)^2 T_Y^2)}{2(\alpha+\beta)} F_y,$$

so that $\deg(\pi_B^\star L) = \frac{(c_1(\mathcal{G})^2 - (\alpha+\beta)^2 T_Y^2)}{2(\alpha+\beta)} - \deg(\pi_B^\star M)$ and we get

$$c_2(\mathcal{G}) = \frac{\alpha}{2(\alpha+\beta)} c_1^2 + \frac{\alpha}{2}(\beta - \alpha) T_Y^2 + (\beta - \alpha)\deg(\pi_B^\star M).$$

Finally, observe that as $\mathcal{G}$ is weakly positive, so is the quotient line bundle $\mathcal{O}_Y(\alpha T_Y + \pi_B^\star M)$, hence $\alpha T_Y + \pi_B^\star M$ is nef.

Indeed, as, by definition, for any ample divisor H on Y and for any $N > 0$ the bundle

$$\left(Sym^N \mathcal{O}_Y(\alpha T_Y + \pi_B^\star M)\right) \otimes \mathcal{O}_Y(H) = \mathcal{O}_Y(N(\alpha T_Y + \pi_B^\star M) + H)$$

is ample, the claim follows.

In particular we have $(\alpha T_Y + \pi_B^\star M) \cdot T_Y \ge 0$, hence $\deg(\pi_B^\star M) \ge -\alpha T_Y^2$.

Finally, as $T_Y^2 \leq 0$ by assumption, we get the statement.

In the case $\alpha = \beta$, the claim can be proved in the same way as above, by taking into account that $\pi_B^\star \pi_{B\star}\mathcal{G}(-\beta T_Y)$ has rank two and its Bogomolov discriminant Δ is zero as such a vector bundle is the pull back of a vector bundle on a curve. On the other hand, the exact sequence

$$0 \to \pi_B^\star \pi_{B\star}\mathcal{G}(-\beta T_Y) \to \mathcal{G}(-\beta T_Y) \to \mathcal{I}_Z \to 0$$

gives $Z = \Delta(\mathcal{G}(-\beta T_Y))$, and as Δ is invariant under shifting by a divisor, we get

$$4c_2(\mathcal{G}) - c_1(\mathcal{G})^2 = \Delta(\mathcal{G}) = \Delta(\mathcal{G}(-\beta T_Y)) \geq 0.$$

Remark. This inequality follows also from the fact that Y is a fibration on B and the restriction of $\mathcal{G}$ to the general fiber is Bogomolov semistable, hence $\mathcal{G}$ is Bogomolov semistable by [Mo, Theorem 2.2.1].

□

Remark. The uniformity condition in Theorem 2.8, (2) is open in the two cases in which we apply Theorem 2.8.

Nevertheless, we believe that the uniformity condition could be dropped by using the theory on weakly positive coherent sheaves and possibly the assumption $Z \cap T_Y = \emptyset$.

As a consequence we directly get the following.

Theorem 2.9. *Let $f : S \to B$ be a fibration with general fiber a trigonal curve, which factorizes through a degree three cover $\rho : S \to Y$ of a ruled surface $\pi_B \colon Y \to B$, and let $\mathcal{E} = (\rho_\star \omega_{S/Y})/\mathcal{O}_Y$. Then*

1. *$s(f) \geq \frac{24(g-1)}{5g+1}$ with equality if and only if the ramification divisor R satisfies $R \equiv \frac{1}{3}\rho^\star \rho_\star R$;*
2. *if the general fiber is a general trigonal curve, then*

$$s(f) \geq \begin{cases} 5 - \frac{8}{g+1} & \text{for } g \text{ odd and } \mathcal{E} \text{ uniform;} \\ \\ 5 - \frac{6}{g} = \mathcal{F}_3(g) & \text{for } g \text{ even.} \end{cases}$$

2.2. **The fourgonal case.** Now we treat the case $n = 4$.

Proposition 2.10. *Let $f \colon S \to B$ be a fibration with general fiber a fourgonal curve, and assume that $f \colon S \to B$ factorizes through a Gorenstien degree four cover $\rho : S \to Y$ where Y is a ruled surface over B. Let $\mathcal{F}$ be the rank two vector bundle $\rho_\star \mathcal{I}_{S,\mathbb{P}(\mathcal{E})}(2)$. Then*

$$R^2 = 2c_1(\mathcal{E})^2 - 4c_2(\mathcal{E}) + c_2(\mathcal{F}), \tag{2.3}$$

hence

$$s(f) = \frac{\frac{2(g+1)}{(g+3)}c_1(\mathcal{E})^2 - 4c_2(\mathcal{E}) + c_2(\mathcal{F})}{\frac{(g+2)}{2(g+3)}c_1(\mathcal{E})^2 - c_2(\mathcal{E})}. \tag{2.4}$$

Proof. The push forward of the exact sequence

$$0 \to \mathcal{I}_{S,\mathbb{P}(\mathcal{E})}(2) \to \mathcal{O}_{\mathbb{P}(\mathcal{E})}(2) \to \mathcal{O}_S(2) \to 0$$

gives

$$0 \to \mathcal{F} \to Sym^2(\mathcal{E}) \to \rho_\star \mathcal{O}_S(2R) \to 0.$$

As by Lemma 2.2 $c_1(\rho_\star \mathcal{O}_S(2R)) \equiv 3c_1(\mathcal{E})$ and $c_2(\rho_\star \mathcal{O}_S(2R)) = c_2(\mathcal{E}) + 4c_1(\mathcal{E})^2 - R^2$, and as

$$c_1(Sym^2(\mathcal{E})) = 4c_1(\mathcal{E}), \quad c_2(Sym^2(\mathcal{E})) = 5c_2(\mathcal{E}) + 5c_1(\mathcal{E})^2, \tag{2.5}$$

we get

$$c_1(\mathcal{F}) = c_1(\mathcal{E}), \qquad c_2(\mathcal{F}) = R^2 + 4c_2(\mathcal{E}) - 2c_1(\mathcal{E})^2.$$

The slope equality follows from Proposition 2.4. □

Now by a detailed study of the ideal sheaf of the surface S inside $\mathbb{P}(\mathcal{E})$ we are able to bound $c_2(\mathcal{F})$ and $c_2(\mathcal{E})$ in terms of $c_1(\mathcal{E})^2$. Finally this will give some bounds on the slope. A first estimate on $c_2(\mathcal{E})$ in terms of $c_2(\mathcal{F})$ and $c_1(\mathcal{E})^2$ follows from Lemma 2.5:

Corollary 2.11. *We have*

$$c_2(\mathcal{E}) \geq \frac{1}{4}\Big(c_1(\mathcal{E})^2 + c_2(\mathcal{F})\Big). \tag{2.6}$$

Proof. Lemma 2.5 implies $c_1(\mathcal{E})^2 \geq R^2 = 2c_1(\mathcal{E})^2 - 4c_2(\mathcal{E}) + c_2(\mathcal{F})$ and the claim follows. □

Corollary 2.12. *Let $f\colon S \to B$ be a fibration, which factorizes through a degree four cover $\rho : S \to Y$, where $\pi_B : Y \to B$ is a ruled surface over B. Then*

$$s(f) \geq 4 + \frac{c_2(\mathcal{F}) - \frac{2}{(g+3)}c_1(\mathcal{E})^2}{\frac{(g+1)}{4(g+3)}c_1(\mathcal{E})^2 - \frac{1}{4}c_2(\mathcal{F})}. \tag{2.7}$$

Proof. As the expression for the slope in Proposition 2.10 is an increasing function in $c_2(\mathcal{E})$, the claim follows from Corollary 2.11.

□

Finally we bound $c_2(\mathcal{F})$. We need to recall a famous result by Schreyer [Sch] on fourgonal curves.

Theorem 2.13. *The canonical model of any fourgonal curve C of genus $g \geq 5$ is a complete intersection of two divisors of the type $2H - b_1\mathbb{P}_y$ and $2H - b_2\mathbb{P}_y$ in its three-dimensional gonal scroll W, where H is the tautological divisor, $\mathbb{P}_y$ is a fiber of W, and the integers b_1 and b_2 satisfy*

$$0 \leq b_2 \leq b_1 \leq g-5, \quad b_1 + b_2 = g-5.$$

Moreover, the degree four covering $C \to \mathbb{P}^1$ induced by the g^1_4 factorizes through double cover of a curve of genus $\gamma < \frac{(g-3)}{6}$ if and only if $b_1 > \frac{2}{3}(g-3)$ and $\gamma = \frac{b_2}{2} + 1$.

We present now our bounds. We denote by C_y the fiber of $f\colon S \to B$ and by F_y the fiber of $\pi_B\colon Y \to B$ over the point $y \in B$, and by $\rho_y\colon C_y \to F_y \cong \mathbb{P}^1$ the restriction of the degree four cover.

Theorem 2.14. *Let $f\colon S \to B$ be a fibration, which factorizes through a degree four cover $\rho : S \to Y$, where $\pi_B : Y \to B$ is a ruled surface over B. Then if the general C_y is a general fourgonal curve of odd genus g, the rank 2 vector bundle $\mathcal{F} = p_\star\mathcal{I}_{S,\mathbb{P}}(2)$ satisfies*

$$c_2(\mathcal{F}) \geq \frac{1}{4}c_1(\mathcal{E})^2. \tag{2.8}$$

Moreover, if the general fiber C_y is any fourgonal curve and $\mathcal{F}$ is weakly positive and uniform on Y, we have

1. *$c_2(\mathcal{F}) \geq \frac{2}{(g+3)}c_1(\mathcal{E})^2$;*
2. *$c_2(\mathcal{F}) \geq \frac{1}{6}c_1(\mathcal{E})^2$ if for the general $y \in B$, $\rho_y\colon C_y \to \mathbb{P}^1$ does not factorize;*
3. *$c_2(\mathcal{F}) \geq \dfrac{(g+2)}{4(g+3)}c_1(\mathcal{E})^2$ if the general C_y is a general fourgonal curve of even genus g.*

Proof. Let (α, β), $\alpha \leq \beta$ be the splitting type of $\mathcal{F}$ on the general fiber of Y. We are going to apply Theorem 2.8 to the rank two vector bundle $\mathcal{F}$.

We need to bound α. Consider a general fiber $C_y \subset S$ and set $V_y = \mathbb{P}(\mathcal{E} \otimes \mathcal{O}_{F_y})$. By definition we have $\mathcal{F} \otimes \mathcal{O}_{F_y} = \rho_\star\mathcal{I}_{S,\mathbb{P}}(2T_\mathbb{P}) \otimes \mathcal{O}_{F_y}$. By projection formula $\rho_\star\mathcal{I}_{S,\mathbb{P}}(2T_\mathbb{P}) \otimes \mathcal{O}_{F_y} \cong \rho_\star(\mathcal{I}_{S,\mathbb{P}}(2T_\mathbb{P}) \otimes \rho^\star\mathcal{O}_{F_y})$, so

$$\mathcal{F} \otimes \mathcal{O}_{F_y} = \rho_\star(\mathcal{I}_{C_y,V_y}(2T_{\mathbb{P}|V_y})).$$

since $\mathcal{I}_{C_y,V_y}(2T_{\mathbb{P}})_{|V_y} = \mathcal{I}_{S,\mathbb{P}}(2T_{\mathbb{P}}) \otimes \rho^\star\mathcal{O}_{F_y}$.

By Theorem 2.13 we can study the minimal free resolution of $\mathcal{I}_{C_y,V_y}$. Indeed, let $W_y \subset \mathbb{P}^{g-1}$ be the scroll containing the canonical model of the general fiber C_y. Let H_y be the hyperplane section of $W_y \subset \mathbb{P}^{g-1}$. A minimal free resolution of $\mathcal{I}_{C_y,W_y}(2H_y)$ has the form:

$$0 \to \mathcal{O}_{W_y}(-2H_y + (g-5)\mathbb{P}_y) \to \mathcal{O}_{W_y}(b_1\mathbb{P}_y) \oplus \mathcal{O}_{W_y}(b_2\mathbb{P}_y) \to \mathcal{I}_{C_y,W_y}(2H_y) \to 0.$$

Recall also that if $g \geq 10$, the g^1_4 on the general fiber is unique, so the two dimensional scroll containing a four-gonal curve as a four-secant curve is unique. As S embeds in $\mathbb{P}$ by the linear system determined by the ramification divisor R, the vector bundle giving the gonal scroll $W_y = \mathbb{P}(\mathcal{H}_y)$, which contains the canonical model of C_y, satisfies

$$\mathcal{E} \otimes \mathcal{O}_{F_y} \cong \mathcal{H}_y(-K_{\mathbb{P}^1}).$$

Now consider the isomorphism between the two scrolls:

$$\phi : V_y \subset \mathbb{P} \to W_y \subset \mathbb{P}^{g-1},$$

and set $T_{\mathbb{P}|V_y} =: T_y$. By definition it follows that $H_y \sim \phi^\star(T_y) - 2\mathbb{P}_y$. Then a minimal free resolution of $\mathcal{I}_{C_y,V_y}(2T_y)$ is given by:

$$0 \to \mathcal{O}_{W_y}(-2T_y + (g+3)\mathbb{P}_y) \to \mathcal{O}_{W_y}((b_1+4)\mathbb{P}_y) \oplus \mathcal{O}_{W_y}((b_2+4)\mathbb{P}_y) \to \mathcal{I}_{C_y,V_y}(2T_y) \to 0.$$

$$\rho_\star\mathcal{O}_{W_y}(-2T_y + (g+3)\mathbb{P}_y) = R^1\rho_\star\mathcal{O}_{W_y}(-2T_y + (g+3)\mathbb{P}_y) = 0,$$

we have

$$\rho_\star\mathcal{I}_{C_y,V_y}(2T_y) \cong \mathcal{O}_{\mathbb{P}^1}(b_1+4) \oplus \mathcal{O}_{\mathbb{P}^1}(b_2+4),$$

so

$$\alpha = b_2 + 4, \quad \beta = b_1 + 4, \quad \alpha + \beta = g + 3.$$

By Theorem 2.13 we have $\alpha \geq 4$, and if the cover $\rho_{C_y} : C_y \to \mathbb{P}^1$ for C_y does not factorize, $\alpha \geq \frac{(g+3)}{3}$. Taking into account Theorem 2.8, this proves claims (1) and (2).

Finally, assume that the general fiber of $f : S \to B$ is a *general* fourgonal curve. From the stratification of the fourgonal locus $\overline{\mathcal{M}}^1_{g,4}$ given in [BSa], it follows that

$$\begin{array}{ll} \alpha = \beta = \frac{(g+3)}{2} & \text{if g is odd,} \\ \alpha = \beta - 1 = \frac{(g+2)}{2} & \text{if g is even.} \end{array}$$

Again, taking into account the inequality in Theorem 2.8, the claimed bounds (2.8) and (3) follow. □

Remark. The weak positivity and uniformity assumptions are fulfilled in the following example: let $\pi_B \colon Y \to B$ be a ruled surface on a smooth curve B, let T_Y be a very ample section of π_B and let $\mathcal{E}$ be a very ample rank three vector bundle on Y. Set $\pi : \mathbb{P}(\mathcal{E}) \to Y$ to be the natural projection and H the tautological divisor on $\mathbb{P}(\mathcal{E})$. Finally, let $S \subset \mathbb{P}(\mathcal{E})$ be the complete intersection surface of two very ample divisors $2H - \alpha\pi^\star T_Y$ and $2H - \beta\pi^\star T_Y$ with $\beta \geq \alpha \geq 1$. Then the vector bundle $\mathcal{F} \cong \mathcal{O}_Y(\alpha T_Y) \oplus \mathcal{O}_Y(\beta T_Y)$ in this case, so it is very ample and uniform.

As a consequence we have the following result.

Theorem 2.15. *Let S be a normal surface. Let Y be a ruled surface over a smooth curve B and let $\pi_B \colon Y \to B$ be the morphism induced by the ruling. If $f : S \to B$ is a fibration which factorizes through a Gorenstein degree four cover $\rho : S \to Y$, then*

(1) if the general fiber C_y of $f : S \to B$ is a general fourgonal curve of odd genus g, then

$$s(f) \geq \frac{16}{3} - \frac{16}{3(3g+1)}.$$

With the further assumptions $\mathcal{F}$ weakly positive and uniform on Y, we have:

(2) $s(f) \geq 4$; if the equality holds, the gonal degree four covering $\rho_{C_y} : C_y \to \mathbb{P}^1$ for C_y a general fiber of $f : S \to B$ factorizes through a double cover of an elliptic curve;

(3) if the 4-cover $\rho_{C_y} : C_y \to \mathbb{P}^1$ for C_y for a general fiber factorizes through a double cover of a curve of genus $\gamma < \frac{(g-3)}{6}$, then

$$s(f) \geq 4 + 4\frac{(\gamma-1)}{(g-\gamma)};$$

(4) if the gonal degree four covering $\rho_{C_y} : C_y \to \mathbb{P}^1$ for C_y a general fiber of $f : S \to B$ does not factorize, then

$$s(f) \geq \frac{24(g-1)}{(5g+3)},$$

with equality if and only if $R \equiv \frac{1}{4}\rho^\star\rho_\star R$;

(5) if the general fiber C_y of $f : S \to B$ is a general fourgonal curve of even genus g, then

$$s(f) \geq \frac{16}{3} - \frac{8}{g} = \mathcal{F}_4(g).$$

In particular the slope conjecture holds for $n = 4$ under our hypotheses.

Proof. This is a straight consequence of Lemma 2.5, Theorem 2.8, Theorem 2.14, Proposition 2.7 and the fact that such a bound is an increasing function in $c_2(\mathcal{F})$. □

Remark.

3. SLOPE OF GENERAL SEMISTABLE FIBRATIONS IN TRIGONAL AND FOURGONAL CURVES.

Recall that if the general fiber is an n - gonal non-hyperelliptic curve of genus $g > (n-1)^2$, so that the g^1_n is unique, then, up to a base change, we may glue the g^1_n's over the fibers to form a linear system on S. This linear system gives a *generically finite rational map* over a ruled surface. Moreover we can choose S to be of general type with birational canonical map and such that the canonical map restricts to the canonical map on a fiber.

This construction relies on base changes, which do not affect the slope. The following result is well-known to the experts (see, for instance, [HMo]).

Theorem 3.1. *Let $f_1 \colon S_1 \to B_1$ be a non isotrivial semistable fibration with fiber of genus $g > 2$ obtained by normalization and base change of a curve $B \subset \overline{\mathcal{M}}_g$ via semistable reduction. Assume that the general fiber is a smooth n-gonal curve and that $g > (n-1)^2$. Then there exists a base change $\overline{B} \to B_1$ such that the if $f \colon S \to \overline{B}$ is the fibration obtained by the base change then $s(f) = s(f_1)$. Moreover there exists a ruled surface $\pi_B \colon \mathbb{P}(\mathcal{B}) \to \overline{B}$ and a rational map $\rho \colon S \dashrightarrow \mathbb{P}(\mathcal{B})$ such that f is factorized through ρ.*

Moreover, the canonical map of S is a birational map and $|K_C| = |K_S|_C$ where C is a fiber of $f \colon S \to \overline{B}$.

By Theorem 3.1, in order to evaluate the slope of a non isotrivial semistable fibration with fiber an n-gonal curve it is sufficient to study the following diagram:

$$\begin{tikzcd} X \arrow[r, "\pi"] \arrow[d, "\tau"'] & \mathbb{P}(\mathcal{B}) \arrow[d, "\pi_B"] \\ S \arrow[r, "f"] & \overline{B}, \end{tikzcd} \tag{3.1}$$

where S is a minimal surface of general type, in particular K_S is big and nef, $\tau\colon X \to S$ is the minimal resolution of $\rho\colon S \dashrightarrow \mathbb{P}(\mathcal{B})$. As in the finite case, we shall still denote by Y the ruled surface $\mathbb{P}(\mathcal{B})$.

A similar construction has already been illustrated by Harris and Morrison in [HMo]. In order to estimate the slope, they preferred to use an intermediate between the family of admissible covers and the semistable model $S \to \overline{B}$. We rely on their analysis. In their case the ruled surface Y is a product surface $\overline{B} \times \mathbb{P}^1$. They blow-up $A \to Y$ in order to have a finite cover $G \to A$ where G dominates S: see [HMo, Diagram (2.2), page 338] (observe that in [HMo] Y and X denote two curves, our S corresponds to their F, our $\overline{B}$ corresponds to their Y). By the very explicit local analytic description of the cover $G \to A$ and of the blow-up $A \to \overline{B} \times \mathbb{P}^1$ done in [HMo, from page 343 to top of page 347], it follows that S differs from G over points of Y, where the branch divisor of the morphism $G \to \overline{B} \times \mathbb{P}^1$ acquires a simple node.

We stress that in a neighbourhood of the so-called points of type (3) [HMo, pages 344 - 346], the rational map $S \dashrightarrow Y = \overline{B} \times \mathbb{P}^1$ is a *finite* morphism, even if the branch locus has not simple nodes [HMo, page 345]. We denote by $B(\pi)$ the branch locus of $\pi\colon X \to Y$. From now on we assume that the followig condition holds:

($\star$) the morphism $\pi\colon X \to Y$ is a finite morphism over a neighbourhood of any point of $B(\pi)$, which is not a simple node.

Definition 3.2. An HM-curve is a smooth curve $B \subset \overline{\mathcal{M}}^1_{g,n}$ such that the condition ($\star$) holds.

Let $\overline{\mathcal{M}}^1_{g,3}$, respectively $\overline{\mathcal{M}}^1_{g,4}$, denote the closure in $\overline{\mathcal{M}}_g$ of the trigonal locus, respectively the closure of the fourgonal locus. Assume that $B \subset \overline{\mathcal{M}}^1_{g,3}$ or $B \subset \overline{\mathcal{M}}^1_{g,4}$ is an HM-curve. By Section 2 of [HMo] the union of such curves is definitely Zariski dense in $\overline{\mathcal{M}}^1_{g,3}$ or $\overline{\mathcal{M}}^1_{g,4}$, so such curves furnish a good test on the slope.

We point out that by Definition 3.2, over singular points of $B(\pi)$ which are not simple nodes the surface S obtained by semistable reduction coincides with the surface X and X coincides with the Stein factorization of $\pi\colon X \to Y$. In particular, a *general* curve in the class constructed in [HMo] is a HM-curve and in the notation of [HMo] over such points, A coincides with $\overline{B} \times \mathbb{P}^1$ and G coincides with S.

We are going to prove that the finite case bounds on the slope hold also for HM-curves. The outline of the proof is the following. We consider the Stein factorization of $\pi\colon X \to Y$, in order to get a finite flat morphism $h : \widehat{X} \to Y$ between a normal surface $\widehat{X}$ and the ruled surface Y.

Now it is possible to construct a Gorenstein model $\widetilde{X}$ and a Gorenstein finite cover $\widetilde{\rho}\colon \widetilde{X} \to \widetilde{Y}$ of a blow up of Y, and to determine precisely the invariants, just because by the condition ($\star$) we need to analyse only germs of non Gorenstein cover of degree 3 or 4 branched over a germ of a simple node.

Finally, we express the slope in terms of the Chern classes of the vector bundles $\widetilde{\mathcal{E}}$, arising from $\widetilde{\rho}_\star\mathcal{O}_{\widetilde{X}} \cong \mathcal{O}_{\widetilde{Y}} \oplus \widetilde{\mathcal{E}}^\vee$, and $\widetilde{\mathcal{F}} = (\widetilde{\pi}_{\mathbb{P}(\widetilde{\mathcal{E}})})_\star(\mathcal{I}_{\widetilde{X},\mathbb{P}(\widetilde{\mathcal{E}})}(2))$, and we bound the second Chern classes of such bundles by using similar arguments to those found in the finite case.

We can conclude by the fact that S is a minimal surface, and if $\widetilde{f}:\widetilde{X}\to\overline{B}$ denotes the induced fibration on the Gorenstein model $\widetilde{X}$, the relation $s(f)\geq s(\widetilde{f})$ holds.

3.1. **The trigonal case.** The following results concern the birational Gorenstein models $\widetilde{X}$ of $\widehat{X}$ in the degree three case, and the numerical class of the branch divisor, which is necessary to compute the invariants of $\widetilde{f}:\widetilde{X}\to\overline{B}$.

Theorem 3.3. *Let X be a normal surface and let $\pi\colon X\to Y$ be a flat degree three morphism to a ruled surface Y. If the non-Gorenstein points of X lay over simple nodes of $B(\pi)$, then there is a finite number of blow-ups $\sigma:\widetilde{Y}\to Y$ and a partial resolution $\widetilde{\tau}:\widetilde{X}\to X$ of X, such that $\widetilde{X}$ is Gorenstein and the induced degree three morphism $\widetilde{\rho}:\widetilde{X}\to\widetilde{Y}$ is Gorenstein.*

Moreover, the possible non-Gorenstein points of X are total ramification points, and

$$\widetilde{\rho}_\star\widetilde{R}\equiv 2\left((g+2)T_0+\frac{c_1^2(\widetilde{\mathcal{E}})+4t}{2(g+2)}F_y-2\sum_{j=1}^{t}E_j\right),$$

where $T_0=T_{\widetilde{Y}}-\frac{1}{2}T^2_{\widetilde{Y}}F_y$ with $T_{\widetilde{Y}}$ any section of $\widetilde{Y}$, F_y is a fiber of $\widetilde{Y}$, the E_j are (-1)-curves arising from the non-Gorenstein points and $\widetilde{\mathcal{E}}$ is determined by $\widetilde{\rho}_\star\mathcal{O}_{\widetilde{X}}\cong\mathcal{O}_{\widetilde{Y}}\oplus\widetilde{\mathcal{E}}^\vee$.

Proof. Let P be a node of $B(\pi)$. We need to study the local behaviour of the covering $\pi\colon X\to Y$ over an analytical neighboorhood of P isomorphic to the polydisk $\Delta\subset\mathbb{C}^2$. We will follow [L]. We know that if $\pi\colon X\to Y$ is a finite and dominant morphism from a variety X with isolated singularities to a smooth variety Y and in Y there is a simple normal crossing divisor D such that π is smooth over $Y\backslash D$ then X has algebraic abelian quotient singularities. In our case to analyse the behaviour on P we can locally identify $B(\pi)\subset Y$ with $\{xy=0\}\subset\mathbb{C}^2_{x,y}$ and P with $(0,0)$. The local fundamental group of $\mathbb{C}^2_{x,y}\setminus\{xy=0\}$ near the origin is free abelian of rank 2, generated by loops around the two boundary divisors. Thus π is analitically equivalent to a covering of the form $\mathbb{C}^2_{u,v}\setminus\{uv=0\}\to\mathbb{C}^2_{x,y}\setminus\{xy=0\}$ determined by the corresponding morphism of fundamental groups. Since the degree is 3 we have to consider subgroups of index ≤ 3.

Explicitly, let $N=\mathbb{Z}^2$ be the lattice of 1-parameter subgroups of the torus $(\mathbb{C}^*)^2_{x,y}$. Let $N'\subset N$ be a sublattice of index 2. Then we easily find that $\widetilde{X}$ is Gorenstein above P.

Let $N'\subset N$ be a sublattice of index 3. This means that over the node we have an open set of X which is generically mapped 3-to-1. Then the inclusion $N'\subset N$ corresponds to a degree three toric morphism $Z\to\mathbb{C}^2$ ramified along the toric boundary divisors. Since π has degree three, the only non Gorenstein possibility is given by $Z=1/3(1,1)=\mathbb{C}^2_{u,v}/(\mathbb{Z}/3\mathbb{Z})$. To find the resolution of such a non-Gorenstein point Q denote by and $E_Q\subset\widetilde{X}$ the exceptional divisor over Q. It follows that E_Q is an irreducible reduced rational curve $E_Q^2=-3$, $E_Q\to E$ is a 3-to-1 cover of the exceptional curve $E\subset\widetilde{Y}$ given by blowing-up P. Hence $K_{\widetilde{X}}=\tau^\star K_X-\frac{1}{3}E_Q$. Then by standard projection formula $E_Q\cdot\widetilde{R}=4$. Let E_i be the exceptional curves on $\widetilde{Y}$ where $\widetilde{\rho}^{-1}(E_i)=E_{Q_i}$, $i=1,...,t$ are the (-3)-exceptional curves we have introduced to obtain our Gorenstein model $\widetilde{X}$. We may express

$$c_1(\widetilde{\mathcal{E}})=(g+2)T_0+kF_y+\sum_{i=1}^{t}a_iE_i$$

and letting $c_1(\widetilde{\mathcal{E}})^2 =: c_1^2$, we get $k = \frac{c_1^2+\sum a_i^2}{2(g+2)}$. Since $\widetilde{\rho}^\star E_i = E_{Q_i}$ and $\widetilde{\rho}_\star \widetilde{R} \equiv 2c_1(\widetilde{\mathcal{E}})$ by Lemma 2.2 , it follows that

$$4 = \widetilde{R} \cdot E_{Q_k} = \widetilde{R} \cdot \widetilde{\rho}^\star E_k = \widetilde{\rho}_\star \widetilde{R} \cdot E_k = 2\left((g+2)T_0 + \frac{c_1^2 + \sum a_i^2}{2(g+2)} F_y + \sum a_i E_i \right) \cdot E_k = -2a_k,$$

hence $a_k = -2$ for any $k = 1, \dots, t$.

□

Theorem 3.4. *Let $\pi \colon X \to Y$ be a generically finite degree three covering between a normal surface X and a ruled surface $\pi_B \colon Y \to B$ over a smooth curve B such that $f_X \colon X \to B$ is a fibration where $f_X = \pi_B \circ \pi$. Assume that the Stein factorization of $\pi \colon X \to Y$ satisfies the hypotheses of Theorem 3.3.*

Then the statements in Theorem 2.9 hold for the slope $s(\widetilde{f})$ of the induced fibration $\widetilde{f} \colon \widetilde{X} \to B$ on the Gorenstein model given by Theorem 3.3.

Proof. Let $\widetilde{\rho} : \widetilde{X} \to \widetilde{Y}$ be the Gorenstein cover constructed in Theorem 3.3. We have

$$c_1(\widetilde{\mathcal{E}}) \equiv (g+2)T_0 + \frac{c_1^2(\widetilde{\mathcal{E}}) + 4t}{2(g+2)} F_y - 2\sum_{j=1}^{t} E_j, \quad K_{\widetilde{Y}} \equiv -2T_0 + \pi^\star K_B + \sum_{j=1}^{t} E_j,$$

so

$$c_1(\widetilde{\mathcal{E}}) \cdot K_{\widetilde{Y}} = 2(b-1)(g+2) - \frac{c_1(\widetilde{\mathcal{E}})^2}{(g+2)} + 2\frac{g}{(g+2)}t, \qquad K^2_{\widetilde{Y}} = -8(b-1) - t.$$

Applying the formula $\widetilde{R}^2 = 2c_1(\widetilde{\mathcal{E}})^2 - 3c_2(\widetilde{\mathcal{E}})$ we get

$$K^2_{\widetilde{f}} = \frac{2g}{(g+2)} c_1(\widetilde{\mathcal{E}})^2 - 3c_2(\widetilde{\mathcal{E}}),$$

$$\chi_{\widetilde{f}} = \frac{(g+1)}{2(g+2)} c_1(\widetilde{\mathcal{E}})^2 - c_2(\widetilde{\mathcal{E}}) + \frac{g}{(g+2)}t.$$

Next we observe that the latter is an increasing function in $c_2(\widetilde{\mathcal{E}})$, so we need to bound it in terms of $c_1(\widetilde{\mathcal{E}})^2$ from below. By similar arguments used in Proposition 2.7 and Theorem 2.8, one can prove the following:

Proposition 3.5. *The second Chern class of the rank 2 vector bundle $\widetilde{\mathcal{E}}$ satisfies:*

(1) $c_2(\widetilde{\mathcal{E}}) \geq \frac{2}{9} c_1(\widetilde{\mathcal{E}})^2$;

(2) $c_2(\widetilde{\mathcal{E}}) \geq \frac{(g+1)}{(g+2)} \left(\frac{1}{4} c_1(\widetilde{\mathcal{E}})^2 + t \right)$ *if the general fiber of $\widetilde{f} : \widetilde{X} \to B$ is a general trigonal curve of odd genus g and $\widetilde{\mathcal{E}}$ is uniform on the proper transforms of the fibers of Y in the blow-up $\widetilde{Y} \to Y$;*

(3) $c_2(\widetilde{\mathcal{E}}) \geq \frac{1}{4} c_1(\widetilde{\mathcal{E}})^2 + t$ *if the general fiber of $\widetilde{f} \colon \widetilde{X} \to B$ is a general trigonal curve of even genus g.*

Proof. The first inequality follows from $\widetilde{R}^2 = 2c_1(\widetilde{\mathcal{E}})^2 - 3c_2(\widetilde{\mathcal{E}})$ and Lemma 2.5.

The other inequalities follow from

$$c_2(\widetilde{\mathcal{E}}) \geq \frac{\alpha}{2(g+2)} \left(c_1(\widetilde{\mathcal{E}})^2 + 4t \right),$$

where (α, β), $\alpha \leq \beta$, is the splitting type of $\widetilde{\mathcal{E}}$ on the general fiber of $\widetilde{Y}$, which can be proved just as in the case of a Gorenstein cover of a ruled surface; see: Theorem 2.8.

We observe that in this case the uniformity condition on the fibres of Y is replaced by the uniformity condition on the proper transforms of the fibers of Y, as the zero-dimensional scheme Z

arising in Brosius sequence does not intersect the exceptional locus in $\widetilde{Y}$. Indeed, in the odd genus case, such a sequence has the form

$$0 \to \mathcal{O}_{\widetilde{Y}}\left(\frac{(g+3)}{2}T_{\widetilde{Y}} + \widetilde{\pi}_B^\star L\right) \to \widetilde{\mathcal{E}} \to \mathcal{O}_{\widetilde{Y}}\left(\frac{(g+1)}{2}T_{\widetilde{Y}} + \widetilde{\pi}_B^\star M - 2\sum_{j=1}^{t} E_j\right) \otimes \mathcal{I}_Z \to 0.$$

If we restrict it to an exceptional divisor E, we get

$$0 \to \mathcal{O}_E \to \widetilde{\mathcal{E}} \otimes \mathcal{O}_E \to \mathcal{O}_E(2) \otimes \mathcal{I}_Z \to 0.$$

On the other hand, $c_1(\widetilde{\mathcal{E}} \otimes \mathcal{O}_E) = c_1(\widetilde{\mathcal{E}}) \cdot E = 2$, so $Z \cap E = \emptyset$.

In the even genus case the Brosius sequence looks like

$$0 \to \widetilde{\pi}_B^\star(\widetilde{\pi}_B)_\star \widetilde{\mathcal{E}}\left(-\frac{(g+2)}{2}T_{\widetilde{Y}}\right) \to \widetilde{\mathcal{E}}\left(-\frac{(g+2)}{2}T_{\widetilde{Y}}\right) \to \mathcal{I}_Z \to 0,$$

so that $c_2(\widetilde{\mathcal{E}}(-\frac{(g+2)}{2}T_{\widetilde{Y}})) = c_2(\widetilde{\mathcal{E}}) + c_1(\widetilde{\mathcal{E}}) \cdot (-\frac{(g+2)}{2}T_{\widetilde{Y}}) + \left(\frac{(g+2)}{2}\right)^2 T_{\widetilde{Y}}^2 = \deg\ Z \geq 0$ and the statement follows. ☐

Remark. In the even genus case $\widetilde{\mathcal{E}}$ turns out to be semistable by [Mo], as in the case of a covering of a ruled surface. Hence we could use the Bogomolov inequality. We prefer to use the better bound $c_2(\widetilde{\mathcal{E}}) \geq \frac{1}{4}c_1(\widetilde{\mathcal{E}})^2 + t$ arising from Brosius sequence, which takes into account the particular expression found for $c_1(\widetilde{\mathcal{E}})$, as this allows to conclude the proof of Theorem 3.4.

To conclude the proof of Theorem 3.4 it is enough to subsitute in the expression

$$s(\widetilde{f}) = \frac{\frac{2g}{(g+2)}c_1(\widetilde{\mathcal{E}})^2 - 3c_2(\widetilde{\mathcal{E}})}{\frac{(g+1)}{2(g+2)}c_1(\widetilde{\mathcal{E}})^2 - c_2(\widetilde{\mathcal{E}}) + \frac{g}{(g+2)}t}$$

the bounds found in Proposition 3.5 and to notice that the so obtained functions are all bounded from below by the same expressions evaluated in $t = 0$. ☐

3.2. **The fourgonal case.**

Theorem 3.6. *Let X be a normal surface and let $\pi\colon X \to Y$ be a flat degree four morphism to a ruled surface Y. If the non-Gorenstein points of X lay over simple nodes of $B(\pi)$, then there is a finite number of blow-ups $\sigma : \widetilde{Y} \to Y$ and a partial resolution $\widetilde{\tau} : \widetilde{X} \to X$ of X, such that $\widetilde{X}$ is Gorenstein and the induced degree four morphism $\widetilde{\rho} : \widetilde{X} \to \widetilde{Y}$ is Gorenstein.*

Moreover, the possible non Gorenstein points of X are total ramification points or ramification points of index three of π, and

$$\widetilde{\rho}_\star \widetilde{R} \equiv 2\left((g+3)T_0 + \frac{c_1^2(\widetilde{\mathcal{E}}) + 9s + 4t}{2(g+3)}F_y - 3\sum_{i=1}^{s} E_i' - 2\sum_{j=1}^{t} E_j''\right),$$

where $\widetilde{R}$ is the ramification divisor of $\widetilde{\rho}$, $T_0 = T_{\widetilde{Y}} - \frac{1}{2}T_{\widetilde{Y}}^2 F_y$ with $T_{\widetilde{Y}}$ any section of $\widetilde{Y}$, F_y is a fiber of $\widetilde{Y}$, E_i' are (-1)-curves arising from total ramification non-Gorenstein points and E_j'' are (-1)-curves arising from index 3 non-Gorenstein points.

Proof. The argument is similar to the one in the proof of Theorem 3.3.

Let P be a node of $B(\pi)$. Since the degree is 4 we have to consider subgroups of index ≤ 4 of. $N = \mathbb{Z}^2$.

Let $N' \subset N$ be a sublattice of index 4. Then the inclusion $N' \subset N$ corresponds to a degree 4 toric morphism $Z \to \mathbb{C}^2$ ramified along the toric boundary divisors. Since the degree is 4 the only case where Z is not Gorenstein is given by $Z = 1/4(1,1) = \mathbb{C}^2_{u,v}/(\mathbb{Z}/4\mathbb{Z})$ with $\mathbb{Z}/4\mathbb{Z}$-action $(u,v) \mapsto (iu, iv)$, and $Z \to \mathbb{C}^2$ given by $(u,v) \mapsto (x,y) = (u^4, v^4)$. Now if $\widetilde{Z} \to Z$ is the minimal resolution and $\widetilde{Y} \to \mathbb{C}^2$ is the blowup of P, then $\widetilde{Z} \to \widetilde{Y}$ is finite flat. By the resolution of quotient singularities $1/4(1,1)$ we construct the claimed partial resolution $\tau\colon \widetilde{X} \to X$ for such points.

Let $Q_1, \dots Q_s \in X$ be the non-Gorenstein singular points of total ramification and let E_{Q_i} denote the corresponding exceptional divisors on $\widetilde{X}$. Set $E'_i := \widetilde{\rho}(E_{Q_i})$ for $i = 1, \dots, s$. Then it holds that

- $2K_{\widetilde{X}} = \tau^\star(2K_X) - \sum_{i=1}^s E_{Q_i}$ where each E_{Q_i} is a rational curve;
- $E_{Q_i}^2 = -4$ and $\widetilde{\rho} : E_{Q_i} \to E$ is $4:1$ on E_{Q_i}, $i = 1, \dots, s$;
- $E_Q \cdot \widetilde{R} = 6$ and $E_Q \not\subset \widetilde{R}$, where $\widetilde{R}$ denotes the ramification divisor of $\widetilde{\rho}$.

We have $K_{\widetilde{X}} = \tau^\star K_X + aE_Q$ and $-2 = E_Q^2 + aE_Q^2 = -4 - 4a$, hence $a = -\frac{1}{2}$.

Now we perform the analysis of a sublattice $N' \subset N$ of index 3. This means that over the node we have an open set of X which is mapped 1-to-1 and another which is generically mapped 3-to-1. The only non-Gorenstein possibility is given by $Z = 1/3(1,1) = \mathbb{C}^2_{u,v}/(\mathbb{Z}/3\mathbb{Z})$. Denote by $E_T \subset \widetilde{X}$ the exceptional divisor over a non-Gorenstein point T of this kind. It follows that E_T is an irreducible reduced rational curve, $E_T^2 = -3$, $E_T \to E''$ is a 3-to-1 cover of the exceptional curve $E'' \subset \widetilde{Y}$ given by blowing-up P. Moreover $E_T \cdot \widetilde{R} = 4$. Notice that to resolve the morphism we have to add a -1-curve $A_T \subset \widetilde{X}$ which is disjoint from E_T and which is mapped 1-to-1 over $E'' \subset \widetilde{Y}$. Notice that $\widetilde{R} \cdot A_T = 0$ but if we contract A_T we can't factorise to $\widetilde{Y}$. We shall denote by $E''_1, \dots, E''_t$ all the exceptional curves on $\widetilde{Y}$ of this type.

Finally observe that we may express

$$c_1(\widetilde{\mathcal{E}}) \equiv \frac{1}{2}\widetilde{\rho}_\star \widetilde{R} \equiv (g+3)T_0 + dF_y + \sum_{i=1}^s a_i E'_i + \sum_{j=1}^t b_j E''_j.$$

Set $c_1^2 := c_1(\widetilde{\mathcal{E}})^2$. We get

$$d = \frac{c_1^2 + \sum a_i^2 + \sum b_j^2}{2(g+3)}.$$

Moreover

$$\widetilde{\rho}^\star E'_i = E_{Q_i}, \quad \widetilde{\rho}^\star E''_j = E_{T_j} + A_{T_j}, \quad \widetilde{\rho}_\star \widetilde{R} \equiv 2c_1(\widetilde{\mathcal{E}}),$$

where $T_1, \dots, T_t$ are the index three non-Gorenstein points.

It follows that

$$6 = \widetilde{R} \cdot E_{Q_k} = \widetilde{R} \cdot \widetilde{\rho}^\star E'_k = \widetilde{\rho}_\star \widetilde{R} \cdot E'_k =$$

$$= 2\left((g+3)T_0 + \frac{c_1^2 + \sum_{i=1}^s a_i^2 + \sum_{j=1}^t b_j^2}{2(g+3)} F_y + \sum_{i=1}^s a_i E'_i + \sum_{j=1}^t b_j E''_j\right) \cdot E'_k = -2a_k,$$

hence $a_k = -3$ for any $k = 1, \dots, s$. Similarly we find

$$4 = \widetilde{R} \cdot (E_{T_l} + E'_{T_l}) = \widetilde{R} \cdot \widetilde{\rho}^\star E''_l = \widetilde{\rho}_\star \widetilde{R} \cdot E''_l =$$

$$= 2\left((g+3)T_0 + \frac{c_1^2 + \sum_{i=1}^s a_i^2 + \sum_{j=1}^t b_j^2}{2(g+3)} F_y + \sum_{i=1}^s a_i E'_i + \sum_{j=1}^t b_j E''_j\right) \cdot E''_l = -2b_l,$$

hence $b_l = -2$ for any $l = 1, \dots, t$.

□

Theorem 3.7. *Let $\pi\colon X \to Y$ be a generically finite degree four covering between a normal surface X and a ruled surface $\pi_B\colon Y \to B$ over a smooth curve B such that $f_X\colon X \to B$ is a fibration where $f_X = \pi_B \circ \pi$. Assume that the Stein factorization of $\pi\colon X \to Y$ satisfies the hypotheses of Theorem 3.6.*

Then the statements in Theorem 2.15 hold for the slope $s(\widetilde{f})$ of the induced fibration $\widetilde{f}\colon \widetilde{X} \to B$ on the Gorenstein model given by Theorem 3.6.

Proof. We have $\widetilde{R}^2 = 2c_1(\widetilde{\mathcal{E}})^2 - 4c_2(\widetilde{\mathcal{E}}) + c_2(\widetilde{\mathcal{F}})$, where $\widetilde{\mathcal{F}} = (\widetilde{\pi}_{\mathbb{P}})_\star \mathcal{I}_{\widetilde{X},\mathbb{P}(\widetilde{\mathcal{E}})}(2)$, and $\chi_{\widetilde{f}} = \chi(\mathcal{O}_{\widetilde{X}}) - (g-1)(b-1) = 4\chi(\mathcal{O}_{\widetilde{Y}}) + \frac{1}{2}c_1(\widetilde{\mathcal{E}})\cdot K_{\widetilde{Y}} + \frac{1}{2}c_1(\widetilde{\mathcal{E}})^2 - c_2(\widetilde{\mathcal{E}}) - (g-1)(b-1)$. Now recall that

$$c_1(\widetilde{\mathcal{E}}) \equiv (g+3)T_0 + \frac{c_1^2(\widetilde{\mathcal{E}}) + 9s + 4t}{2(g+3)}F_y - 3\sum_{i=1}^{s} E_i - 2\sum_{j=1}^{t} E''_j, \quad K_{\widetilde{Y}} \equiv -2T_0 + \pi^\star K_B + \sum_{i=1}^{s} E'_i + \sum_{j=1}^{t} E''_j,$$

hence

$$c_1(\widetilde{\mathcal{E}})\cdot K_{\widetilde{Y}} = 2(b-1)(g+3) - \frac{c_1(\widetilde{\mathcal{E}})^2}{(g+3)} + 3\frac{g}{(g+3)}s + 2\frac{(g+1)}{(g+3)}t, \qquad K^2_{\widetilde{Y}} = -8(b-1) - s - t.$$

It follows that

$$K^2_{\widetilde{f}} = 2\frac{(g+1)}{(g+3)}c_1(\widetilde{\mathcal{E}})^2 - 4c_2(\widetilde{\mathcal{E}}) + c_2(\widetilde{\mathcal{F}}),$$

$$\chi_{\widetilde{f}} = \frac{(g+2)}{2(g+3)}c_1(\widetilde{\mathcal{E}})^2 - c_2(\widetilde{\mathcal{E}}) + \frac{3g}{2(g+3)}s + \frac{(g+1)}{(g+3)}t,$$

As $\widetilde{X}$ is normal, we may apply Lemma 2.5 and get

$$c_2(\widetilde{\mathcal{E}}) \geq \frac{1}{4}\Big(c_1(\widetilde{\mathcal{E}})^2 + c_2(\widetilde{\mathcal{F}})\Big).$$

These relations give

$$s(\widetilde{f}) \geq 4 + \frac{c_2(\widetilde{\mathcal{F}}) - \frac{2}{(g+3)}c_1(\widetilde{\mathcal{E}})^2}{\frac{(g+1)}{4(g+3)}c_1(\widetilde{\mathcal{E}})^2 - \frac{1}{4}c_2(\widetilde{\mathcal{F}}) + \frac{3g}{2(g+3)}s + \frac{(g+1)}{(g+3)}t}. \tag{3.2}$$

Next we observe that the latter is an increasing function in $c_2(\widetilde{\mathcal{F}})$, and the $c_2(\widetilde{\mathcal{F}})$ can be bounded using similar arguments as in the ruled case. The statement will follow as the resulting expressions are all bounded from below by the same expressions evaluated in $s = t = 0$. □

We point out that in the following Proposition in the general odd genus case $\widetilde{\mathcal{F}}$ turns out to be Bogomolov semistable, but we prefer to consider the Brosius sequence also in this case, as it gives a better bound on the $c_2(\widetilde{\mathcal{F}})$ than the Bogomolov inequality.

More precisely we have:

Proposition 3.8. *The second Chern class of $\mathcal{F}$ satisfies:*

(1) $c_2(\widetilde{\mathcal{F}}) \geq \frac{1}{4}\left(c_1(\widetilde{\mathcal{E}})^2 + 9s + 4t\right)$ *if the general fiber C_y of $\widetilde{f} : \widetilde{X} \to B$ is a general fourgonal curve of odd genus g;*

If moreover $\widetilde{\mathcal{F}}$ is weakly positive and uniform on the proper transforms of the fibers of Y in the blow-up $\widetilde{Y} \to Y$, then:

(2) $c_2(\widetilde{\mathcal{F}}) \geq \frac{2}{(g+3)}\left(c_1(\widetilde{\mathcal{E}})^2 + 9s + 4t\right)$*;*

(3) $c_2(\widetilde{\mathcal{F}}) \geq \frac{1}{6}\left(c_1(\widetilde{\mathcal{E}})^2 + 9s + 4t\right)$ *if the gonal degree four covering* $\rho_{C_y} : C_y \to \mathbb{P}^1$ *for* C_y *a general fiber of* $\widetilde{f} : \widetilde{X} \to B$ *does not factorize;*

(4) $c_2(\widetilde{\mathcal{F}}) \geq \dfrac{(g+2)}{4(g+3)}\left(c_1(\widetilde{\mathcal{E}})^2 + 9s + 4t\right)$ *if the general fiber* C_y *of* $\widetilde{f} : \widetilde{X} \to B$ *is a general fourgonal curve of even genus* g*.*

Proof. The proof is similar to the one in the trigonal case of Proposition 3.5, taking into account the bounds on β given in the proof of Theorem 2.14. □

We can finally conclude with our main result.

Theorem 3.9. *Let* $B \subset \overline{\mathcal{M}}^1_{g,3}$*, with* $g \geq 5$*, or* $B \subset \overline{\mathcal{M}}^1_{g,4}$*, with* $g \geq 10$*, be an HM-curve. Let* $f : S \to \overline{B}$ *be a semistable reduction obtained from* B*.*

Then the statements in Theorems 2.9 and 2.15 hold for f*; in particular the Harris-Stankova conjecture holds for a HM-curve if* $n = 4$ *and the genus is odd, and it holds for a general (not intersecting a codimension two subvariety) HM-curve if* $n = 4$*, the genus is even and the vector bundle* $\widetilde{\mathcal{F}}$ *is weakly positive.*

Proof. Fix a curve B as in the statement; after normalizing $B_1 \to B$ and by a base change $\overline{B} \to B_1$ we obtain a semistable fibration $f\colon S \to \overline{B}$ as in Theorem 3.1. We know that $s(f) = 12 - s(B)$. Let $\pi\colon X \to Y$ be the generically finite morphism naturally associated to the n-gonal fibration $f\colon S \to \overline{B}$. Let $h\colon \widehat{X} \to Y$ be the finite morphism arising from the Stein factorization of π. The surface $\widehat{X}$ is a normal one; see: Lemma 3.10 below.

Now we recall that a finite morphism between a normal and a smooth surface is flat, hence $h\colon \widehat{X} \to Y$ is a finite and flat morphism.

By condition $(\star)$ the possible non Gorenstein points of $\widehat{X}$ can spring out of germs of Y where the branch locus has a simple node. Then the claims follow from Theorem 3.4 and Theorem 3.7.

□

Only for lack of reference we prove this well-known result:

Lemma 3.10. *Let* X *and* Y *be two smooth schemes. Let* $\pi\colon X \to Y$ *be a generically finite morphism. Let* $\pi\colon X \xrightarrow{g} \widehat{X} \xrightarrow{h} Y$ *be the Stein factorization of* $\pi\colon X \to Y$*. Then* $\widehat{X}$ *is normal.*

Proof. The problem is local since we have to show that for every point $z \in \widehat{S}$ its local ring $\mathcal{O}_{\widehat{S},z}$ is integrally closed in $\mathrm{Frac}(\mathcal{O}_{\widehat{S},z}) = \mathbb{C}(\widehat{S}) = \mathbb{C}(X)$. By [Ha, Corollary 11.5 and Corollary 11.3] it holds that $g_\star\mathcal{O}_X = \mathcal{O}_{\widehat{S}}$. By definition $\mathcal{O}_{\widehat{S},z} = \underline{\lim}_{z\in U}\mathcal{O}_{\widehat{S}}(U) = \underline{\lim}_{z\in U}\mathcal{O}_X(g^{-1}U)$. Since X is smooth then $\mathcal{O}_X(g^{-1}U)$ is normal, that is it is integrally closed in $\mathbb{C}(X)$, since $g^{-1}(U)$ is an open set of the topological space associated to the scheme X. Now we are reduced to show that the direct limit of a family of normal subrings of a field is a normal ring. Let $\{A\}_{i\in I}$ be a directed family of integrally closed subrings of a field K. Let $A = \underline{\lim}_{i\in I}A_i$. Let $\alpha \in K$ and $n \in \mathbb{N}$ such that $\alpha^n + a_1\alpha^{n-1} + \ldots + a_{n-1}\alpha + a_0 = 0$ where $a_i \in A$, $i = 0, 1, \ldots, n-1$. By definition of direct limit, there exists an $m \in I$ such that there exist $b_i \in A_m$ such that $\alpha^n + b_1\alpha^{n-1} + \ldots + b_{n-1}\alpha + b_0 = 0$. Since A_m is normal then $\alpha \in A_m$. Then $\alpha \in A$.

□

REFERENCES

[AW] M. Artin, G. Winters, *Degenerate fibres and stable reduction of curves*, Topology 10 (1971), 373 - 383.

[B] M. A. Barja, *On the slope and geography of fibred surfaces and threefolds*, Ph. D. Thesis, University of Barcelona, 1998.

[BS] M.A. Barja, L. Stoppino, *Slopes of trigonal fibred surfaces and of higher dimensional fibrations*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **8** (2009), no. 4, 647-658.

[BZ] M. A. Barja and F. Zucconi, *On the slope of fibred surfaces*, Nagoya Math. J. **164** (2001), 103-131.

[B] J.E. Brosius, *Rank-2 vector bundles on a ruled surface. I.*, Math. Ann. **265** (1983), no. 2, 155 - 168.

[BSa] M. Brundu, G. Sacchiero, *Stratification of the moduli space of fourgonal curves*, preprint (2012), arXiv:1204.6156v1.

[CE] G. Casnati, T. Ekedahl, *Covers of algebraic varieties. I. A general structure theorem, covers of degree 3,4 and Enriques surfaces*, J. Algebraic Geom. 5 (1996), no. 3, 439 - 460.

[Ca] F. Catanese, *Fibred Surfaces, Varieties Isogenous To a Product and related Moduli Spaces*, American Journal of Mathematics Vol. **122**, No. 1 (2000), pp. 1 - 44.

[CFM] D. Chen, G. Farkas, I. Morrison, *Effective divisors on moduli spaces of curves*, preprint (2012), arXiv:1205.6138v1.

[CH] M. Cornalba, J. Harris, *Divisor classes associated to families of stable varieties, with applications to the moduli space of curves.* Ann. Sc. Ec. Norm. Sup. **21 (4)** (1988), 455-475.

[CS] M. Cornalba and L. Stoppino, *A sharp bound for the slope of double cover fibrations*, Michigan Math. J. 56 (2008), no. 3, 551-561.

[EH] D. Eisenbud and J. Harris, *The Kodaira dimension of the moduli space of curves of genus* ≥ 23, Invent. Math. 90 (1987), 359-387.

[Fa] G. Farkas, *The geometry of the moduli space of curves of genus 23,* Math. Ann. 318 (2000), 43-65.

[FP] G. Farkas and M. Popa, *Effective divisors on* M_g*, curves on K3 surfaces, and the slope conjecture*. J. Algebraic Geom. 14 (2005), no. 2, 241-267.

[FJ] M. Fedorchuk, D. Jensen, *Stability of 2nd Hilbert points of canonical curves,* preprint (2011), arXiv:1111.5339v1.

[Fr] R. Friedman, *Algebraic surfaces and holomorphic vector bundles,* Universitext, Springer-Verlag, New York, 1998. 328 pp.

[HMo] J. Harris and I. Morrison, *Slopes of effective divisors on the moduli space of stable curves*, Invent. Math. 99 (1990), 321-355.

[HM] J. Harris and D. Mumford, *On the Kodaira dimension of the moduli space of curves*, Invent. Math. 67 (1982), 23-86.

[Ha] R. Hartshorne, *Algebraic geometry*, Grad. Texts Math., Vol. 52, Springer Verlag, New York-Heidelberg-Berlin, (1977).

[K1] K. Konno, *A lower bound of the slope of trigonal fibrations*, Internat. J. Math. vol. **7** no. 1 (1996), 19-27.

[K2] K. Konno, *Clifford index and the slope of fibreed surfaces,* J. Algebraic Geom. 8 n.2 (1999), 207-220.

[L] H. Laufer, *Normal two dimensional singularities*, Ann. of Math. studies 71, Princeton University Press, (1971).

[Mo] A. Moriwaki, *A sharp slope inequality for general stable fibrations of curves*, J. Reine Angew. Math. **480** (1996), 177-195.

[M] D. Mumford, *Stability of Projective Varieties*, L'Ens. Math. 23 (1977) 39-110.

[P] R. Pandharipande *Descendent bounds for effective divisors on* $\overline{\mathcal{M}}_g$, J. Algebraic Geometry, vol. **21**, (2012), 299-303.

[Sch] F. O. Schreyer, *Syzygies of canonical curves and special linear series*, Math. Ann. 275 (1986), no. 1, 105137.

[S] Z. Stankova, *Moduli of trigonal curves,* J. Algebraic Geom. 9 (2000), no. 4, 607-662.

[St] L. Stoppino, *Slope inequalities for fibred surfaces via GIT*, Osaka J. Math. 45 (2008), no. 4, 1027-1041.

[Sz] L. Szpiro, *Séminaire sur les pinceaux de courbes de genre au moins deux*, Astérisque 86 (1981), 142.

[V] E. Viehweg, *Die Additivitt der Kodaira Dimension fr projektive Faserrume ber Varietten des allgemeinen Typs*, Journ. reine angew. Math. **330** (1982), 132 - 142

[X1] G. Xiao, *Irregularity of surfaces with a linear pencil*, Duke Math. J. vol. **55**, no. 3 (1987), 597-602.

[X2] G. Xiao, *Fibred algebraic surfaces with low slope.* Math. Ann. **276** (1987), 449-466.

Dipartimento di Matematica e Geoscienze, Università degli studi di Trieste,, via Valerio 12/b, 34127 Trieste, Italy, beorchia@units.it

Dipartimento di Matematica e Informatica, via delle Scienze 206, Università degli studi di Udine, Udine, 33100 Italy, Francesco.Zucconi@uniud.it